

Day : Wednesday

Date: 6/23/2004

Time: 11:00:49

PALM INTRANET

# Inventor Information for 10/043658

Inventor Name	City	State/Country
OLSON, ERIC N.	DALLAS	TEXAS

[Appln Info](#)[Contents](#)[Petition Info](#)[Atty/Agent Info](#)[Continuity Data](#)[Foreign Data](#)Search Another: Application# or Patent# PCT /  /  or PG PUBS # Attorney Docket #  Bar Code #  

To go back use Back button on your browser toolbar.

Back to [PALM](#) | [ASSIGNMENT](#) | [OASIS](#) | [Home page](#)

Day : Wednesday

Date: 6/23/2004

Time: 11:00:55

**PALM INTRANET**

## Continuity Information for 10/043658

### Parent Data

10043658is a continuation of 09438075Which Claims Priority from Provisional Application 60107755Which Claims Priority from Provisional Application 60108083

### Child Data

No Child Data

Appln Info	Contents	Petition Info	Atty/Agent Info	Continuity Data	Foreign Data	Invent
------------	----------	---------------	-----------------	--------------------	--------------	--------

Search Another: Application# or Patent# PCT /  / or PG PUBS # Attorney Docket # Bar Code # 

To go back use Back button on your browser toolbar.

Back to [PALM](#) | [ASSIGNMENT](#) | [OASIS](#) | [Home page](#)



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Br

Search PubMed for mef2 heart hypertrophy

Limits Preview/Index History Clipboard Details

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Show:

Items 1-7 of 7

One page

☐ 1: [Czubryt MP, Olson EN.](#)

Related Articles, Link:



Balancing contractility and energy production: the role of myocyte enhancer factor 2 (MEF2) in cardiac hypertrophy.

Recent Prog Horm Res. 2004;59:105-24. Review.

PMID: 14749499 [PubMed - indexed for MEDLINE]

☐ 2: [Zhang CL, McKinsey TA, Chang S, Antos CL, Hill JA, Olson EN.](#)

Related Articles, Link:



Class II histone deacetylases act as signal-responsive repressors of cardiac hypertrophy.

Cell. 2002 Aug 23;110(4):479-88.

PMID: 12202037 [PubMed - indexed for MEDLINE]

☐ 3: [Liu ZP, Nakagawa O, Nakagawa M, Yanagisawa H, Passier R, Richardson JA, Srivastava D, Olson EN.](#)

Related Articles, Link:



CHAMP, a novel cardiac-specific helicase regulated by MEF2C.

Dev Biol. 2001 Jun 15;234(2):497-509.

PMID: 11397016 [PubMed - indexed for MEDLINE]

☐ 4: [Han J, Molkentin JD.](#)

Related Articles, Link:



Regulation of MEF2 by p38 MAPK and its implication in cardiomyocyte biology.

Trends Cardiovasc Med. 2000 Jan;10(1):19-22. Review.

PMID: 11150724 [PubMed - indexed for MEDLINE]

☐ 5: [Olson EN, Williams RS.](#)

Related Articles, Link:



Remodeling muscles with calcineurin.

Bioessays. 2000 Jun;22(6):510-9. Review. Erratum in: Bioessays 2000 Nov;22(11):1049.

PMID: 10842305 [PubMed - indexed for MEDLINE]

☐ 6: [Passier R, Zeng H, Frey N, Naya FJ, Nicol RL, McKinsey TA, Overbeek P, Richardson JA, Grant SR, Olson EN.](#)

Related Articles, Link:



CaM kinase signaling induces cardiac hypertrophy and activates the MEF2 transcription factor in vivo.

J Clin Invest. 2000 May;105(10):1395-406.

PMID: 10811847 [PubMed - indexed for MEDLINE]

☐ 7: [Kolodziejczyk SM, Wang L, Balazsi K, DeRepentigny Y, Kothary R, Megeney LA.](#)

Related Articles, Link:



MEF2 is upregulated during cardiac hypertrophy and is required for normal post-natal growth of the myocardium.

Curr Biol. 1999 Oct 21;9(20):1203-6.

PMID: 10531040 [PubMed - indexed for MEDLINE]

Show:

Items 1-7 of 7

One page

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Jun 7 2004 18:11



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Br  
 Search PubMed for Go Clear  
 Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

Entrez PubMed  
 Overview  
 Help | FAQ  
 Tutorial  
 New/Noteworthy  
 E-Utilities

PubMed Services  
 Journals Database  
 MeSH Database  
 Single Citation Matcher  
 Batch Citation Matcher  
 Clinical Queries  
 LinkOut  
 Cubby

Related Resources  
 Order Documents  
 NLM Gateway  
 TOXNET  
 Consumer Health  
 Clinical Alerts  
 ClinicalTrials.gov  
 PubMed Central

☐ 1: Recent Prog Horm Res. 2004;59:105-24.

[Related Articles, Link](#)

Full text article at  
[rphr.endojournals.org](http://rphr.endojournals.org)

## Balancing contractility and energy production: the role of myocyte enhancer factor 2 (MEF2) in cardiac hypertrophy.

Czubryt MP, Olson EN.

University of Texas Southwestern Medical Center at Dallas, Dallas, Texas 75390-9148, USA.

Cardiac hypertrophy -- that is, enlargement of the heart resulting from increased myocyte size -- is observed with many forms of human heart disease. It may arise secondary to an insult, such as infarct or chronic hypertension, or may occur as a consequence of a genetic defect, such as in hypertrophic cardiomyopathy. Traditionally, it has been widely believed that hypertrophy occurred as an adaptive response to normalize increased wall stress due to disease. Recently, however, it has been observed that while hypertrophy initially appears to improve the function of the heart following insult, over time, it frequently leads to a decompensated state, characterized by fibrosis and chamber dilation, resulting in overt heart failure. Hypertrophy also occurs during fetal development, immediately after birth, and in trained athletes; however, it does not lead to decompensation in these states. Experiments over the last 15 years have implicated similar signaling pathways in both pathological and physiological hypertrophic responses. Recently, important differences have been demonstrated that might hold the key to the development of effective new treatments for human diseases. This chapter focuses on how these hypertrophic responses differ from one another phenotypically and discusses how inefficient or impaired energy metabolism in the heart may contribute to the development of pathological responses. We also discuss recent evidence that the myocyte enhancer factor 2 (MEF2) transcription factor family, which previously has been shown to be important in cardiac development and hypertrophy, may have a role in regulation of cardiac energy metabolism.

Publication Types:

- Review
- Review, Tutorial

PMID: 14749499 [PubMed - indexed for MEDLINE]

---

Display Abstract Show: 20 Sort Send to Text

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Jun 7 2004 18:11: